













FRIEND

Philippe Berio

Yves Bresson, Jean-Michel Clausse, Denis Mourard, Julien Dejonghe, Stéphane Lagarde, Anthony Meilland, Karine Perraut, Isabelle Tallon-Bosc, Nicolas Nardetto, Alain Spang, Christophe Bailet, Aurélie Marcotto, Philippe Stee, Michel Tallon and CHARA staff

OVERVIEW













- Why FRIEND?
- What is FRIEND?
- FRIEND in the CHARA focal lab
- First Technical Run
- Conclusions and Perspectives

Why FRIEND?











Preparation of the next generation of visible interferometric instruments

- Successor of VEGA, currently in operation at CHARA
- VEGA main limitations

Photon-counting detectors (ALGOL): saturation at high flux and photon centroiding hole Multi-mode regime: limitation in magnitude and accuracy

- ⇒ no low visibility and closure phase measurements
- Installation of AO systems on CHARA (in progress) and on VLTI/AT (NAOMI)
- Very low noise (<0.5e-) and fast (up to 2000 fr/s) analogic detector: OCAM2
- Combination of up to 6 telescopes simultaneously in the visible

FRIEND \Rightarrow validation of spectrally-resolved interferometric observations in the visible in the case of partial correction by AO

What is FRIEND?















- 3 telescopes
- Spatial filtering with mono mode optical fibers (as AMBER and MIRC)
- Photometric channels
- · Multi-axial 'all-in-one' beam recombination scheme
- Dispersed fringes mode (as VEGA, AMBER and MATISSE)
- 2 Spectral Resolution (R=400/ $\Delta\lambda$ =120nm and R=2500/ $\Delta\lambda$ =30nm)
- Use of analogic very low noise camera: OCAM2





Visible Interferometry Workshop January 15-16, <u>20</u>15

FRIEND in the CHARA focal lab










BEAM COMBINATION LABORATORY



FRIEND can use the alignment (pupils and images) and source devices of VEGA.

FRIEND in the CHARA focal lab
















FRIEND Injection Module



Visible Interferometry Workshop January 15-16, 2015


















Visible Interferometry Workshop January 15-1<mark>6, 2015</mark>




















Visible Interferometry Workshop January 15-16, 2015




















FRIEND in the CHARA focal lab














FRIEND Combiner Module



Visible Interferometry Workshop January 15-16, 2015

















Visible Interferometry Workshop January 15-1<mark>6, 2015</mark> Nice


















- 3 Nights allocated
 December 18-19-20, 2014
- FRIEND installed and aligned in 2 days
 Fringes with the VEGA internal source
- Poor seeing and bad weather conditions
 r_o<5cm and Clouds/Humidity
- However first fringes obtained on Regulus 2T and Low Resolution mode
































Observations log

Date	Star	DIT	Gain	Spectral Resolution	Telescopes	Comments
18/12/14	Regulus	10	471	None	E2	Photometric test
18/12/14	Aldebaran	10	471	None	E2	Photometric test
18/12/14	Aldebaran	10	1000	LR	E2	Photometric test
18/12/14	γ Ori	10	1000	None	E2	Photometric test
18/12/14	κ <u>O</u> ri	10	1000	None	E2	Photometric test
18/12/14	Aldebaran	10	471	None	E2	Photometric test
18/12/14	Aldebaran	10	471	None	E2	Photometric test
18/12/14	Aldebaran	100	653	LR	E2	Photometric test
19/12/14	Regulus	40	997	LR	E1E2	Fringes
19/12/14	Regulus	20	997	LR	E1E2	Fringes
19/12/14	Regulus	10	997	LR	E1E2	Fringes
19/12/14	Regulus	5	997	LR	E1E2	Fringes
19/12/14	Regulus	2	997	LR	E1E2	Fringes
19/12/14	γ <u>Cas</u>	20	997	LR	E1E2	?
20/12/14	а Сер	10	997	LR	S1S2	Fringes
20/12/14	κ Ori	10	997	LR	S1S2	?
20/12/14	Regulus	10	997	LR	E1E2	?













Preliminary results

Artifical Source





Why Low visibility?

- ⇒ Polarization effects
- ⇒ Size of the pinehole of the artificial source

















Preliminary results

Observation of Regulus





Short DIT ⇒ reduce the OPD jitter effect ⇒ High instrumental Visibility but low Flux

What is the optimal configuration of the detector?

Conclusions and Perspectives











- Fringes recorded on Regulus (m_v=1.4) and α Cep (m_v=2.4) in 2T/LR mode
- Estimated coupling ~ 0.2 % (bad seeing conditions)
 - ⇒ however consistent with simulation
 - ⇒ expected coupling in case of good seeing ~ 3-5 %
- Fringes Search and Coherencing done with VEGA and CLIMB
- FRIEND does not replace VEGA. Both inst. could work alternatively

Conclusions and Perspectives













- ⇒ 3T observations
- ⇒ Medium spectral resolution
- ⇒ Polarization analysis
- ⇒ Optimal configuration of OCAM2
- ⇒ Low visibility and Closure Phase measurements
- First Scientific Run in September 2015
 - ⇒ Science case to be defined
- Final Goal = Test FRIEND with AO
 - \Rightarrow in 2016?